

# Effects of logging, hunting, and forest fragment size on physiological stress levels of two sympatric ateline primates in Colombia

Rebecca Rimbach<sup>1,2,\*</sup>, Andrés Link<sup>2,3</sup>, Michael Heistermann<sup>4</sup>, Carolina Gómez-Posada<sup>5</sup>, Nelson Galvis<sup>2,3</sup> and Eckhard W. Heymann<sup>1</sup>

<sup>1</sup> Behavioral Ecology and Sociobiology Unit, German Primate Center, Kellnerweg 4, 37077 Göttingen, Germany

<sup>2</sup> Fundación Proyecto Primates, Cra. 11a No. 91-55, Bogotá, Colombia

<sup>3</sup> Departamento de Ciencias Biológicas, Universidad de Los Andes, Cra. 1 No. 18a-12, Bogotá, Colombia

<sup>4</sup> Endocrinology Laboratory, German Primate Center, Kellnerweg 4, 37077 Göttingen, Germany

<sup>5</sup> Department of Biology, University of Washington, 106 Kincaid, Seattle, WA 98195, USA

\*Corresponding author: Behavioral Ecology and Sociobiology Unit, German Primate Center, Kellnerweg 4, 37077 Göttingen, Germany.  
Tel: +49 551 3851 469. Email: rimbach@dpz.eu

Habitat fragmentation and anthropogenic disturbances are of major concern to the conservation of endangered species because of their potentially negative impact on animal populations. Both processes can impose physiological stress (i.e. increased glucocorticoid output) on animals, and chronically elevated stress levels can have detrimental effects on the long-term viability of animal populations. Here, we investigated the effect of fragment size and human impact (logging and hunting pressure) on glucocorticoid levels of two sympatric Neotropical primates, the red howler monkey (*Alouatta seniculus*) and the critically endangered brown spider monkey (*Ateles hybridus*). These two species have been reported to contrast strongly in their ability to cope with anthropogenic disturbances. We collected faecal samples from eight spider monkey groups and 31 howler monkey groups, living in seven and 10 different forest fragments in Colombia, respectively. We measured faecal glucocorticoid metabolite (FGCM) levels in both species using previously validated methods. Surprisingly, fragment size did not influence FGCM levels in either species. Spider monkeys showed elevated FGCMs in fragments with the highest level of human impact, whereas we did not find this effect in howler monkeys. This suggests that the two species differ in their physiological responsiveness to anthropogenic changes, further emphasizing why brown spider monkeys are at higher extinction risk than red howler monkeys. If these anthropogenic disturbances persist in the long term, elevated FGCM levels can potentially lead to a state of chronic stress, which might limit the future viability of populations. We propose that FGCM measurements should be used as a tool to monitor populations living in disturbed areas and to assess the success of conservation strategies, such as corridors connecting forest fragments.

**Key words:** *Alouatta seniculus*, *Ateles hybridus*, faecal glucocorticoid metabolites, habitat fragmentation, hunting, logging

**Editor:** Steven Cooke

Received 28 July 2013; Revised 4 October 2013; Accepted 5 October 2013

**Cite as:** Rimbach R, Link A, Heistermann M, Gómez-Posada C, Galvis N, Heymann EW (2013) Effects of logging, hunting, and forest fragment size on physiological stress levels of two sympatric ateline primates in Colombia. *Conserv Physiol* 1: doi:10.1093/conphys/cot031.

## Introduction

Habitat loss, habitat fragmentation, and anthropogenic disturbances that accompany these processes (e.g. logging, increased hunting pressure) are of major concern to the conservation of endangered species due to their role in population declines (Peres, 2001; Fahrig, 2003). Currently, numerous species from all vertebrate groups are threatened with extinction (IUCN, 2012). The pervasive process of anthropogenic disturbances of natural ecosystems (Hannah *et al.*, 1995; Foley *et al.*, 2005) emphasizes the need to understand the proximate effects that these alterations have on the health and survival of animal populations. Generally, taxa with narrow dietary niches and that occupy few habitat types are at greater extinction risk than taxa that have broader niches and occupy several habitat types (Harcourt *et al.*, 2002). Species respond to fragmentation and disturbances differently depending on factors such as life history, geographical range, ecological niche, and dispersal ability (Purvis *et al.*, 2000; Henle *et al.*, 2004; Cardillo *et al.*, 2005). Some species manage to adjust aspects of their behaviour and social system (Menon and Poirier, 1996; Sumner *et al.*, 1999; González-Solís *et al.*, 2001; Blumstein *et al.*, 2005; Umapathy *et al.*, 2011), whereas species that cannot adjust face local extinction (Cosson *et al.*, 1999; Peres, 2001).

The role that changes in physiological parameters play in population declines and, ultimately, species extinction is not yet well understood. Glucocorticoids (GCs; cortisol and corticosterone, depending on the species) are the frontline hormones of the vertebrate stress response (Sapolsky *et al.*, 2000) and potentially play an important role. Many different stimuli (e.g. predators, aggression from conspecifics, food deprivation) can elicit a stress response that triggers the release of GCs. Such elevations of GC levels are adaptive and important for the survival of the individual (Boonstra, 2005; Monclús *et al.*, 2005). Glucocorticoids mobilize readily available energy that can be used to respond to the stimulus that triggered the release of GCs (Selye, 1956; Breazile, 1987; Stratakis and Chrousos, 1995). During the stress response, other energetically demanding activities, such as digestion and reproduction, are suppressed (Wingfield *et al.*, 1998; Landys *et al.*, 2006). Hence, long-term elevations of GCs can have deleterious effects on reproduction, growth, and immune system activity (Pickering *et al.*, 1991; Cyr and Romero, 2007; Charbonnel *et al.*, 2008; French *et al.*, 2010; Setchell *et al.*, 2010). Anthropogenic disturbances have been associated with GC elevations in many vertebrate taxa (amphibians: Homan *et al.*, 2003; Janin *et al.*, 2011; reptiles: French *et al.*, 2010; birds: Lucas *et al.*, 2006; Wasser *et al.*, 1997; and mammals: Martínez-Mota *et al.*, 2007; Gobush *et al.*, 2008; Rangel-Negrín *et al.*, 2009; Jaimez *et al.*, 2012; Dunn *et al.*, 2013). These GC elevations have been linked to negative effects on reproduction and immune system activity (e.g. Ellenberg *et al.*, 2007; French *et al.*, 2010), suggesting that elevated stress levels caused by human influence can

directly affect individual health and, ultimately, population viability.

We studied two primate species that occur sympatrically in Colombia, namely brown spider monkeys (*Ateles hybridus*) and red howler monkeys (*Alouatta seniculus*), to investigate whether and how anthropogenic disturbances influence their GC output. The two genera *Ateles* and *Alouatta* have been reported to contrast strongly in their ability to cope with anthropogenic disturbances (Bernstein *et al.*, 1976; Bicca-Marques, 2003; Michalski and Peres, 2005). Brown spider monkeys are endemic to Colombia and Venezuela, and are considered to be one of the 25 most endangered primate species worldwide due to severe habitat loss and high hunting pressure (Mittermeier *et al.*, 2012). Habitat fragmentation, slow reproductive cycles, large area requirements, and their dietary niche (for review, see Di Fiore *et al.*, 2010) make this species vulnerable to anthropogenic disturbances. Moreover, spider monkeys exhibit flexible grouping patterns (fission–fusion dynamics) that are likely to reduce intra-group feeding competition and enable them to cope with changes in the availability of ripe fruit, their preferred food resources (e.g. Klein and Klein, 1977; Symington, 1988). Accordingly, the confinement to small fragments might reduce the flexibility of their grouping patterns and thereby lower their potential to minimize competition, which might cause increased physiological stress.

In contrast, red howler monkeys seem less vulnerable to anthropogenic disturbances and are able to persist even in extremely small fragments (Crockett, 1998; Lopez *et al.*, 2005; Michalski and Peres, 2005). They have a broad distribution range, occupy a wide array of ecosystems, and are currently not threatened with extinction (Boubli *et al.*, 2008). Typically, howler monkeys require much smaller home ranges than spider monkeys (reviewed by Di Fiore *et al.*, 2010). They form cohesive groups (Neville, 1972; for review, see Di Fiore and Campbell, 2007) and have a mainly folivorous but flexible diet (Milton, 1980; Julliot and Sabatier, 1993; Bicca-Marques, 2003).

The proximate mechanisms leading to such species-specific differences in the ability to cope with a changing environment have not yet been investigated. Therefore, the aim of this study was to determine how fragment size and level of human impact influence physiological stress levels, measured through faecal glucocorticoid metabolites (FGCMs), in both species. We collected faecal samples from both species in forest fragments that differed in size and level of human impact (hunting and logging activities). We predicted that FGCM levels would increase with both, decreasing fragment size (as a proxy for fragmentation intensity) and increasing level of human impact. Due to differences in diet, area requirements, and adaptation capabilities, we predicted that red howler monkeys would generally react less strongly than brown spider monkeys to these disturbances, and that they would show no elevation in GC levels or a less pronounced elevation compared with brown spider monkeys.

## Materials and methods

### Study sites

Between April 2010 and April 2012, we collected faecal samples from eight spider monkey and 31 howler monkey groups, living in seven and 10 different fragments, respectively. Fragments differed in size and human impact (Table 1). Four fragments (San Juan, Quinchas, Jamaica, and Juntas) are long-term study sites, and study groups have been habituated to human observers previously (Aldana *et al.*, 2008; Gómez-Posada *et al.*, 2010; Link *et al.*, 2010). We visited all other fragments only for faecal sample collection purposes (for 2–3 weeks). Nine of the study fragments were occupied by both species and the other four (Jamaica, Juntas, LGPM, and Cienaga) were occupied only by howler monkeys. Although most fragments contained both species, sometimes faecal sample collection was feasible only for one. We determined the level of current human impact through observations (e.g. recently cut tree stumps, presence of hunting dogs in fragments, pet primates at farms) and surveys in which we interviewed farm owners and workers. We classified fragments without current human hunting activities and absence of recent logging activity ( $\leq 2$  months) as level 0, fragments with either hunting or logging activity as level 1, and fragments with both hunting and ongoing logging activity as level 2.

### Faecal sample collection

We collected samples from adult and subadult individuals (mean  $\pm$  SD: *Ateles*,  $7.9 \pm 12.5$  samples per individual and

*Alouatta*,  $2.3 \pm 2.6$  samples per individual) and, for every sample, noted sex and age-class, female reproductive state (when identifiable), collection time, and date. The time lag of GC metabolite excretion in faeces is  $\sim 24$  h in *A. hybridus* and  $\sim 46$  h in *A. seniculus* (Rimbach *et al.*, 2013). Thus, we avoided following unhabituated groups on 2–3 consecutive days (depending on the species) to reduce the influence of observer presence on FGCM levels. In unhabituated groups, we did not recognize individuals and, to avoid resampling of individuals, we sampled each group or subgroup (in the case of *Ateles*) only once. Faecal samples collected in the fragments ‘Jamaica’ and ‘Juntas’ were collected between May and July 2010, and stored at  $-20^{\circ}\text{C}$  until extraction in December 2012. Long-term faecal sample storage at  $-20^{\circ}\text{C}$  has been shown to preserve immunoreactive faecal GC metabolites reliably (Hunt and Wasser, 2003), and we are therefore confident that this storage method has not affected the FGCM measurements of these samples.

In both species, pregnancy cannot reliably be detected by observation. Thus, to determine the approximate conception date and female reproductive state at sample collection we used parturition date (habituated groups) in combination with average gestation length (*Ateles*:  $\sim 7.5$  months and *Alouatta*:  $\sim 6.3$  months; Di Fiore and Campbell, 2007). We categorized females as lactating for the time in which they nursed dependent offspring. Females that were neither pregnant nor lactating were categorized as non-pregnant, non-lactating. However, in unhabituated groups we were able to categorize females merely as either lactating or non-lactating (pregnant and cycling females). We collected comparative numbers of samples from

**Table 1:** Number of faecal samples collected for each species in the different forest fragments ( $n = 13$ ), which varied in size and level of human impact

Fragment	Size (ha)	Human impact	No. of samples (groups)		Location
			<i>Ateles hybridus</i>	<i>Alouatta seniculus</i>	
San Juan	65	0	411 (2)	289 (17)	6° 43'N, 74° 09'W
San Juan3	75	0	10 (1)	5 (1)	6° 43'N, 74° 07'W
LGPM	100	0	—	9 (1)	6° 41'N, 74° 09'W
Quinchas	250	0	46 (1)	21 (3)	6° 02'N, 74° 16'W
Terra Firme1	500	0	3 (1)	—	6° 41'N, 74° 08'W
LGPM2	500	0	—	4 (1)	6° 41'N, 74° 09'W
Jamaica	4.21	1	—	12 (1)	4° 23'N, 75° 48'W
Juntas	25.5	1	—	13 (1)	4° 25'N, 75° 47'W
India	500	1	3 (1)	—	6° 15'N, 74° 07'W
Cienaga	50	2	—	6 (2)	6° 42'N, 74° 08'W
Campo Capote	250	2	3 (1)	5 (1)	6° 34'N, 73° 51'W
Remedios	400	2	5 (1)	—	6° 53'N, 74° 34'W
San Juan 4	500	2	—	9 (3)	6° 41'N, 74° 07'W
Total no. of samples			481 (8)	373 (31)	

both sexes (*A. hybridus*: 51.8% females and 48.2% males; and *A. seniculus*: 45.6% females and 54.4% males) per fragment and have no reason to assume that the collection was biased towards collecting samples from females in only certain reproductive conditions.

Before placing faecal material into the sample tube, we homogenized the faecal bolus and removed any obvious undigested matter (e.g. large seeds). We placed ~0.5 g of faeces into a 15 ml pre-weighed polypropylene tube pre-filled with 5 ml of 96% ethanol and shook the tube until the faeces were suspended in the solvent (Shutt *et al.*, 2012; Rimbach *et al.*, 2013). We kept the samples at ambient temperatures until steroid extraction in the evening.

### Steroid extraction and faecal glucocorticoid metabolite analysis

We determined faecal wet weight by calculating the difference between the weight of the tube before (tube plus ethanol) and after sample collection (tube, ethanol, and faeces). We shook the tubes firmly for 5 min and thereafter, centrifuged the samples for 1 min using a manually operated centrifuge (Shutt *et al.*, 2012; Rimbach *et al.*, 2013). We poured off ~2 ml of each faecal extract into 2 ml polypropylene tubes, covered them with parafilm, labelled them, and stored them at ambient temperatures (~25°C) in a dark place. Every 2 months we transported the extracts to the Universidad de Los Andes, Bogotá and stored them at -20°C until shipment to the Endocrinology Laboratory at the German Primate Center for analysis. We showed in a previous study that storing faecal extracts in this way did not affect FGCM levels (Rimbach *et al.*, 2013).

We analysed all faecal samples using a previously validated (Rimbach *et al.*, 2013) enzyme-immunoassay for 11 $\beta$ -hydroxyetiocholanolone, a group-specific measurement of 5 $\beta$ -reduced GC metabolites (Ganswindt *et al.*, 2003) with a 3 $\alpha$ ,11 $\beta$ -dihydroxy structure. The enzyme-immunoassay was performed as described in detail by Heistermann *et al.* (2004). Prior to hormone measurement, we diluted extracts 1:250–1:2000 (depending on original concentration) in assay buffer and thereafter, took duplicate aliquots to assay. Intra- and inter-assay coefficients of variation of were 6.1 and 7.8%, respectively, for high-value and 7.4 and 13.0%, respectively, for low-value quality controls. All hormone concentrations are expressed as nanograms per gram faecal wet weight.

### Statistical analyses

To assess the influence of fragment size and level of human impact on FGCM levels, we used linear mixed models (LMM; Baayen, 2010). We fitted all models with the lmer function from the lme4 R-package (Bates and Maechler, 2010) in R2.15.1 (R Development Core Team 2012). We used restricted maximal likelihood methods to estimate the models, because they are robust against unequal sample sizes (Keselman *et al.*, 2001). We used two LMMs per species, and

group and fragment ID were used as random factors in all models. Glucocorticoid excretion often shows a diurnal rhythm, and GC levels can be affected by a variety of potentially confounding factors (for review, see Millsaugh and Washburn, 2004; Keay *et al.*, 2006; Goymann, 2012). To account for these factors, we used sex, age, female reproductive state, group size, and sample collection time as control variables in the models (because some of these variables have been shown to affect FGCM levels in the study species; Rimbach *et al.*, 2013). We tested for interactions between fragment size and human impact, and between group size and fragment size. These were not significant ( $P \geq 0.05$ ) and are not included in the final models. For each study species we used two models, one using the size of each fragment and one comparing fragments with ‘control sites’. In the second set of models, we defined ‘control sites’ as fragments that were larger than a certain, species-specific, size. Due to differences in area requirements between the study species (Di Fiore *et al.*, 2010), we defined fragments larger than 50 ha as ‘control sites’ (or as a proxy for continuous forest) for red howler monkeys, whereas for brown spider monkeys we defined fragments larger than 200 ha as ‘control sites’.

We log transformed the response variable (FGCM levels) to achieve normal distribution and checked that the assumptions of normally distributed and homogeneous residuals were fulfilled by visually inspecting Q–Q plots and the residuals plotted against the fitted values for each model. To assess model stability, we ran diagnostics (dfbetas) that did not suggest the existence of influential cases. We used the function `vif` of the R-package `car` (Fox and Weisberg, 2011) applied to a standard linear model, excluding the random effects, to derive variance inflation factors (Field, 2005). We used a likelihood ratio test (R function ‘`anova`’) to determine the significance of the full model (all fixed and random effects) compared with the corresponding null model (only random effects). We used the functions `pvals.fnc` of the R-package ‘`language R`’ (Baayen, 2010) to determine  $P$  values based on Markov Chain Monte Carlo (MCMC) sampling (Baayen, 2011). All statistical tests were two tailed, and the statistical threshold was set at  $P \leq 0.05$ .

## Results

In both species, FGCM levels varied substantially between different fragments (Figure 1). The fragment size did not affect FGCM levels in spider monkeys ( $n = 481$  samples; LMM:  $\chi^2 = 2.06$ , d.f. = 1,  $P = 0.15$ ; Figure 1a), whereas human impact had an influence (LMM:  $\chi^2 = 17.5$ , d.f. = 2,  $P = 0.0001$ ; Figure 2a). Specifically, spider monkeys living in fragments with both kinds of human influence had FGCM levels (mean  $\pm$  SEM = 496.7  $\pm$  90.0 ng/g) that were more than twice as high as those found in animals living in fragments with no disturbance (mean  $\pm$  SEM = 206.74  $\pm$  7.5 ng/g;  $P_{\text{MCMC}} = 0.01$ ; Table 2) or only one type of disturbance (mean  $\pm$  SEM = 137.16  $\pm$  45.5 ng/g;  $P_{\text{MCMC}} = 0.04$ ). There was no difference between the FGCM levels of spider

monkeys living in fragments with no human impact or only one type ( $P_{\text{MCMC}} = 0.48$ , Table 2).

The results of the second model (comparing fragments with ‘control sites’) were similar to the model using fragment size. The FGCM levels of spider monkeys did not differ between fragments (mean  $\pm$  SEM = 217.28  $\pm$  8.12 ng/g) and ‘control sites’ (mean  $\pm$  SEM = 378.51  $\pm$  106.42 ng/g;  $n = 481$  samples; LMM,  $\chi^2 = 0.27$ , d.f. = 1,  $P = 0.60$ ; Table 3). As in the first model, the level of human impact influenced FGCM


levels of spider monkeys (LMM,  $\chi^2 = 16.21$ , d.f. = 2,  $P = 0.0003$ ; Table 3).

In howler monkeys, neither fragment size (Figure 1b) nor human impact (none: mean  $\pm$  SEM = 245.9  $\pm$  10.6 ng/g; one: 159.4  $\pm$  31.0 ng/g; and both: 251.2  $\pm$  32.9 ng/g; Figure 2b) influenced FGCM levels ( $n = 373$  samples, full vs. null model  $\chi^2 = 13.76$ , d.f. = 9,  $P = 0.13$ ). The second model showed similar results. Neither human impact nor forest type (fragments: mean  $\pm$  SEM = 168.61  $\pm$  26.89 ng/g; and ‘control sites’: mean  $\pm$  SEM = 246.94  $\pm$  10.36 ng/g) influenced FGCM levels of howler monkeys ( $n = 373$  samples, full vs. null model,  $\chi^2 = 11.02$ , d.f. = 9,  $P = 0.27$ ).


### Discussion

Consistent with our predictions, spider monkeys had higher FGCM levels in fragments with the highest level of human impact compared with less impacted fragments. We did not find such an effect in howler monkeys. In contrast to our predictions, neither fragment size nor forest type (fragments compared to ‘control sites’) influenced FGCM levels of either species. Time of sample collection, age, and female reproductive state influenced FGCM levels in *A. hybridus* (possible explanations are discussed elsewhere; see Rimbach *et al.*, 2013), but we controlled for these factors in data analyses. This study reinforces previous results concerning species-specific differences in the ability to cope with anthropogenic disturbances and strengthens the assumption that brown spider monkeys are more susceptible to human alterations than red howler monkeys.

Proximity to humans, hunting, and logging activities are likely to be perceived as threatening by many animals. Red deer (*Cervus elaphus*) that were chased by humans (Bateson and Bradshaw, 1997) and female African elephants (*Loxodonta africana*) that ranged in areas with high poaching risk (Gobush *et al.*, 2008) had elevated GC levels compared to conspecifics that did not experience the disturbance. The presence of humans can lead to elevated GC levels in several animal taxa (birds: Fowler, 1999; Müllner *et al.*, 2004; Thiel *et al.*, 2011;



**Figure 1:** Mean  $\pm$  SEM faecal glucocorticoid metabolite levels of *Ateles hybridus* (a) and *Alouatta seniculus* (b) in relationship to forest fragment size.



**Figure 2:** Mean  $\pm$  SEM faecal glucocorticoid metabolite levels of *Ateles hybridus* (a) and *Alouatta seniculus* (b) in relationship to level of human impact (none: no hunting or logging, one: either hunting or logging, both: both hunting and logging).

**Table 2:** Results of the linear mixed models examining the influence of forest fragment size and level of human impact on log-transformed faecal glucocorticoid metabolite levels in *Ateles hybridus*

Null vs. full model	d.f.	$\chi^2$	P value
	9	113.38	<0.001*
Variable	Estimate ± SEM	t	$P_{MCMC}$
Intercept	5.76 ± 0.33	17.52	0.0001*
Fragment size	0.09 ± 0.05	1.81	0.1694
Impact: both–none	−0.93 ± 0.32	−2.88	0.0134*
Impact: both–one	−1.37 ± 0.57	−2.39	0.0400*
Impact: one–none	0.43 ± 0.59	0.72	0.4820
Group size	0.03 ± 0.04	0.83	0.7428
Time	−0.29 ± 0.03	−8.72	0.0001*
Sex	0.14 ± 0.09	1.48	0.1532
Age	−0.33 ± 0.11	−2.84	0.0062*
Overall effect	d.f.	$\chi^2$	P
Female reproductive state	3	21.15	<0.001*
Fragment size	1	2.06	0.1565*
Human impact	2	17.5	0.0001*

\*Variables that significantly influenced faecal glucocorticoid metabolite levels.

reptiles: French *et al.*, 2010; mammals: Creel *et al.*, 2002; Barja *et al.*, 2007; Behie *et al.*, 2010; Muehlenbein *et al.*, 2012; Piñeiro *et al.*, 2012; Zwijacz-Kozica *et al.*, 2012). Moreover, proximity to humans can impair the breeding success of animals (Ellenberg *et al.*, 2006; Hinam and St Clair, 2008; Strasser and Heath, 2013), which might be caused by increased GC levels (Ellenberg *et al.*, 2007; Charbonnel *et al.*, 2008). Furthermore, logging activities can result in elevated GC levels (Wasser *et al.*, 1997). Concordant with previous studies, we found elevated FGCM levels of brown spider monkeys in fragments where logging and hunting occurred. Whether this elevation of FGCM levels indicates a state of chronic stress with potential negative consequences on health and fitness is difficult to assess, especially in such long-lived animals, and is beyond the scope of this paper (for a discussion of chronic stress, see e.g. Boonstra, 2013). In contrast to spider monkeys, we did not find an effect of logging and hunting on FGCM levels of red howler monkeys, which suggests that this species might have a lower sensitivity to react to anthropogenic disturbances with an activation of the hypothalamic–pituitary–adrenal axis than brown spider monkeys. Likewise, the intensity of human presence (number of tourists) did not influence the physiological stress levels of another howler monkey species (*Alouatta palliata mexicana*; Aguilar-Melo *et al.*, 2013, but see Behie *et al.*, 2010).

Fruit availability often declines in small and established (>10 years) fragments (Putz *et al.*, 1990; Cordeiro and Howe, 2001; Arroyo-Rodríguez and Mandujano, 2006; Dunn *et al.*,

**Table 3:** Results of the linear mixed models examining the influence of forest type (fragment or control sites) and level of human impact on log-transformed faecal glucocorticoid metabolite levels in *Ateles hybridus*

Null vs. full model	d.f.	$\chi^2$	P value
	9	72.52	<0.001*
Variable	Estimate ± SEM	t	$P_{MCMC}$
Intercept	6.33 ± 0.32	19.58	0.0001*
Forest type	−0.20 ± 0.39	−0.531	0.5708
Impact: both–none	−1.06 ± 0.34	−3.05	0.0168*
Impact: both–one	1.04 ± 0.57	1.81	0.0868
Impact: one–none	−1.06 ± 0.34	−3.05	0.0172*
Group size	0.10 ± 0.03	2.61	0.3718
Time	−0.23 ± 0.03	−6.30	0.0001*
Sex	−0.01 ± 0.10	−0.15	0.9004
Age	−0.36 ± 0.12	−2.95	0.0030*
Overall effect	d.f.	$\chi^2$	P value
Female reproductive state	3	8.38	0.0386*
Forest type	1	0.27	0.6033*
Human impact	2	16.21	0.0003*

\*Variables that significantly influenced faecal glucocorticoid metabolite levels.

2010), and low food availability can cause elevated GC levels in primates (Cavigelli, 1999; Muller and Wrangham, 2004; Chapman *et al.*, 2007; Behie *et al.*, 2010). Surprisingly, spider monkeys living in small fragments did not show elevated FGCM levels compared with those living in larger ones, although they potentially experience low levels of food availability and high resource competition. Although some fragments included in this study are very small, drastic changes in food availability might not have occurred yet because most of these fragments have been created rather recently (<10 years). This might explain why fragment size did not influence FGCM levels of either species. Alternatively, it could be that elevated GC levels were associated with low food availability in some fragments, but we are lacking data on resource availability for most fragments and thus, cannot test this assumption. However, two very small fragments (Jamaica and Juntas) were isolated about 100 years ago, and food availability is extremely low (Gómez-Posada *et al.*, 2010). Nevertheless, howler monkeys are able to persist in these fragments and seem to maintain relatively low GC levels.

In contrast to frugivores, folivores are often able to persist in moderately disturbed areas (Johns and Skorupa, 1987), probably because leaf quantity and quality are often higher in disturbed areas, especially at edges, where light exposure is high (Johns, 1988; Ganzhorn, 1995; Irwin, 2008). In the case of howler monkeys, altered leaf availability and quality might compensate for negative effects associated with small fragments

and human impact because of their mainly folivorous diet. This supports the notion that they are capable of habituating to human activities, which is likely to explain why we did not find increased FGCM levels in individuals that live in small and disturbed fragments.

The observed inter-specific differences in responsiveness to human impact could also be the result of a different 'perception' of stressful factors. Differences in population densities between fragments might be a crucial factor determining GC levels of animals, and howler monkeys have been reported to live at very high densities in fragments (Rumiz, 1990; Ostro *et al.*, 2001; Gómez-Posada *et al.*, 2010; Link *et al.*, 2010). Owing to the lack of data on population densities for most fragments, we were not able to include this variable. Future research should include this factor and thus help to determine whether the species shows variation in FGCM levels according to different population density levels.

Generally, our results support previous findings emphasizing a species-specific effect of human disturbance and habitat fragmentation on adrenocortical activity. This specificity might be the reason for inconsistent results revealed by previous studies that used GC levels as markers of physiological stress in a conservation context (reviewed by Busch and Hayward, 2009). While several studies report elevated GC levels in response to anthropogenic disturbances (Wasser *et al.*, 1997; Lucas *et al.*, 2006; Martínez-Mota *et al.*, 2007; Rangel-Negrín *et al.*, 2009; Janin *et al.*, 2011; Jaimez *et al.*, 2012), others found the reverse, or no effect at all. For instance, red-bellied lemurs (*Eulemur rubriventer*; Tecot, 2008) and African forest elephants (*Loxodonta africana cyclotis*; Munshi-South *et al.*, 2008) show higher GC levels in undisturbed areas than conspecifics in disturbed habitats. Canadian grizzly bears (*Ursus arctos*) exhibit lower GC levels in areas with high poaching activity compared with less disturbed areas (Wasser *et al.*, 2004), whereas Alaskan brown bears (*Ursus arctos horribilis*), a closely related subspecies, show no effect of human presence on GCs (von der Ohe *et al.*, 2004). This demonstrates that species probably differ in their sensitivity to disturbances and that not all species respond with a predictable change in GC levels, or (not mutually exclusive) that such a physiological response depends on the degree of the threat perceived.

One important limitation of our study is the lack of faecal samples from extensive and continuous forests and uneven sample sizes between fragments. The small sample size in fragments where primates are being hunted reflects the challenge of encountering and following arboreal animals that are wary and fearful of humans. Although we have only few samples of *A. hybridus* from two fragments with both types of human impact, the FGCM levels of all those samples are much higher than for those collected in other fragments. Thus, it is conceivable that these differences in FGCM levels reflect true differences, although additional studies should be conducted to confirm these results. Another impeding factor for sample collection is the high degree of fragmentation of the remaining

habitat of *A. hybridus* (Urbani *et al.*, 2008) that exacerbated the access to large forests. Our sample size of fragments with only one type of human impact is also small, and future research (with a larger number of fragments) may clarify which of the two factors (logging or hunting) drives the observed elevation in FGCM levels in brown spider monkeys. Moreover, the intensity of human activities is likely to be another important factor influencing FGCM levels. Unfortunately, we were not able to quantify the intensity of hunting and logging activities during this study (e.g. how many people worked in the fragments per day, how many primates were killed in a defined time period). One way to overcome problems concerning the acquisition of data on the intensity of human activities may be to collect samples of the same population before, during, and after logging or hunting activities occur and investigate how FGCM levels change accordingly.

In the absence of data on fitness and health parameters, the results of this study should be interpreted conservatively and they should be confirmed by additional studies. Nevertheless, by controlling for many variables that can potentially confound GC levels (of which some have been neglected previously) and by comparing GC levels of two species that occur (at least partly) in the same fragments, we provide important evidence for species-specific differences in physiological responsiveness and susceptibility to anthropogenic disturbances.

This study reveals that some species (e.g. howler monkeys) may not be negatively influenced by a moderate level of human activity and suggests that agricultural ecosystems could be of use to conserve them. However, our results also demonstrate that GC levels of some species can be elevated in response to anthropogenic disturbances. To what extent these GC elevations reflect a situation of chronic stress with potentially negative fitness consequences or are merely a reflection of an acute adrenocortical reaction to ongoing human activities and, as such, might be adaptive to cope with a short-term challenge (without consequences on fitness) is impossible to assess in the absence of longer-term investigations on fitness and health parameters. It is conceivable, however, that if anthropogenic disturbances persist in the long term, this can potentially lead to a state of chronic stress, which might limit the future viability of animal populations. This study emphasizes the need for the active protection of continuous forests for the conservation of species with low coping abilities (e.g. spider monkeys). Measurements of physiological stress levels should be used to monitor populations living in disturbed areas and to assess the success (concerning amelioration or minimization of stress) of conservation strategies such as corridors connecting fragments and the promotion of alternative sources of animal protein for the human population (e.g. to decrease hunting pressure).

## Acknowledgements

We thank the Lalinde and De Greiff families for permission to work at 'Hacienda San Juan de Carare' and all the persons who

allowed us to visit their farms and forests. We thank Professor Pablo R. Stevenson and the Laboratorio de Ecología de Bosques Tropicales y Primatología for logistic support with sample storage. For their help with sample collection, we thank Felipe Alfonso, Natalia Alvis, Nathalia Fuentes, Miguel García, David Gongora, Claudia Lodoño, Natalia Mejía, Jennifer Rey-Goyeneche, Germán Ríos, and especially Andres Montes. For their support in the laboratory, we acknowledge Andrea Heistermann and Petra Kiesel. We thank Cedric Girard-Buttoz for his statistical advice. Fundación ProAves provided all logistic support for work at Reserva El Paujil (Quinchas). We also thank the Ministerio de Medio Ambiente y Desarrollo Sostenible in Colombia for the permits needed for this study. Fundación Proyecto Primates provided all logistics. The study was conducted completely non-invasively under the research permit 4120-E1-110897. This work was supported by the German Primate Center; Margot Marsh Biodiversity Foundation's Primate Action Fund; Rufford Small Grants Foundation; and Idea Wild.

## References

- Aguilar-Melo AR, Andresen E, Cristóbal-Azkarate J, Arroyo-Rodríguez V, Chavira R, Schondube J, Serio-Silva JC, Cuarón AD (2013) Behavioral and physiological responses to subgroup size and number of people in howler monkeys inhabiting a forest fragment used for nature-based tourism. *Am J Primatol* 75: 1108–1116.
- Aldana AM, Beltrán M, Torres-Neira J, Stevenson PR (2008) Habitat characterization and population density of brown spider monkeys (*Ateles hybridus*) in Magdalena Valley, Colombia. *Neotrop Primates* 15: 46–50.
- Arroyo-Rodríguez V, Mandujano S (2006) Forest fragmentation modifies habitat quality for *Alouatta palliata*. *Int J Primatol* 27: 1079–1096.
- Baayen RH (2010) *Analyzing Linguistic Data: A Practical Introduction to Statistics*. Cambridge University Press, Cambridge, UK.
- Baayen RH (2011) language R: data sets and functions with “Analyzing linguistic data: a practical introduction to statistics”, R package version 1.4. <http://cran.r-project.org/package=languageR>
- Barja I, Silván G, Rosellinic S, Piñero A, González-Gilb A, Camachob L, Illera JC (2007) Stress physiological responses to tourist pressure in a wild population of European pine marten. *J Steroid Biochem Mol Biol* 104: 136–142.
- Bates D, Maechler M (2010) lme4: Linear mixed-effects models using S4 classes. <http://cran.r-project.org/package=lme4>
- Bateson P, Bradshaw EL (1997) Physiological effects of hunting red deer (*Cervus elaphus*). *Proc Biol Sci* 264: 1707–1714.
- Behie AM, Pavelka MSM, Chapman CA (2010) Sources of variation in fecal cortisol levels in howler monkeys in Belize. *Am J Primatol* 72: 600–606.
- Bernstein I, Balcaen P, Dresdale L, Gouzoules H, Kavanagh M, Patterson T, Neyman-Warner P (1976) Differential effects of forest degradation on primate populations. *Primates* 17: 401–411.
- Bicca-Marques JC (2003) How do howler monkeys cope with habitat fragmentation?. In LK Marsh, ed., *Primates in Fragments: Ecology and Conservation*. Kluwer Academic/Plenum Publishers, New York, pp 283–303.
- Blumstein DT, Fernández-Juricic E, Zollner PA, Garity SC (2005) Interspecific variation in avian responses to human disturbance. *J Appl Ecol* 42: 943–953.
- Boonstra R (2005) Equipped for life: the adaptive role of the stress axis in male mammals. *J Mammal* 86: 236–247.
- Boonstra R (2013) Reality as the leading cause of stress: rethinking the impact of chronic stress in nature. *Funct Ecol* 27: 11–12.
- Boubli J-P, Di Fiore A, Rylands AB, Mittermeier RA (2008) *Alouatta seniculus*. In: IUCN 2013. IUCN Red List of Threatened Species. Version 2013.1. <http://www.iucnredlist.org>. (last accessed 25 July 2013).
- Breazile JE (1987) Physiologic basis and consequences of distress in animals. *J Am Vet Med Assoc* 191: 1212–1215.
- Busch DS, Hayward LS (2009) Stress in a conservation context: a discussion of glucocorticoid actions and how levels change with conservation-relevant variables. *Biol Conserv* 142: 2844–2853.
- Cardillo M, Mace GM, Jones KE, Bielby J, Bininda-Emonds ORP, Sechrest W, Orme CDL, Purvis A (2005) Multiple causes of high extinction risk in large mammal species. *Science* 309: 1239–1241.
- Cavigelli SA (1999) Behavioural patterns associated with faecal cortisol levels in free-ranging female ring-tailed lemurs, *Lemur catta*. *Anim Behav* 57: 935–944.
- Chapman CA, Saj TL, Snaith TV (2007) Temporal dynamics of nutrition, parasitism, and stress in colobus monkeys: implications for population regulation and conservation. *Am J Phys Anthropol* 134: 240–250.
- Charbonnel N, Chaval Y, Berthier K, Deter J, Morand S, Palme R, Cosson JF (2008) Stress and demographic decline: a potential effect mediated by impairment of reproduction and immune function in cyclic vole populations. *Physiol Biochem Zool* 81: 63–73.
- Cordeiro NJ, Howe HF (2001) Low recruitment of trees dispersed by animals in African forest fragments. *Conserv Biol* 15: 1733–1741.
- Cosson JF, Ringuet S, Claessens O, de Massary JC, Dalecky A, Villiers JF, Granjon L, Pons JM (1999) Ecological changes in recent land-bridge islands in French Guiana, with emphasis on vertebrate communities. *Biol Conserv* 91: 213–222.
- Creel S, Fox JE, Hardy A, Sands J, Garrott B, Peterson RO (2002) Snowmobile activity and glucocorticoid stress responses in wolves and elk. *Conserv Biol* 16: 809–814.
- Crockett CM (1998) Conservation biology of the genus *Alouatta*. *Int J Primatol* 19: 549–578.
- Cyr NE, Romero ML (2007) Chronic stress in free-living European starlings reduces corticosterone concentrations and reproductive success. *Gen Comp Endocrinol* 151: 82–89.

- Di Fiore A, Campbell CJ (2007) The atelines: variation in ecology, behavior, and social organization. In CJ Campbell, A Fuentes, KC MacKinnon, M Panger, SK Bearder, eds, *Primates in Perspective*. Oxford University Press, New York, pp 155–185.
- Di Fiore A, Link A, Campbell CJ (2010) The atelines: behavioral and socio-ecological diversity in a New World radiation. In CJ Campbell, A Fuentes, KC MacKinnon, M Panger, SK Bearder, eds, *Primates in Perspective*, Ed 2. Oxford University Press, Oxford, pp 155–188.
- Dunn JC, Cristóbal-Azkarate J, Veà JJ (2010) Seasonal variations in the diet and feeding effort of two groups of howlers in different sized forest fragments. *Int J Primatol* 31: 887–903.
- Dunn JC, Cristóbal-Azkarate J, Schulte-Herbrüggen B, Chavira R, Veà JJ (2013) Travel time predicts fecal glucocorticoid levels in free-ranging howlers (*Alouatta palliata*). *Int J Primatol* 34: 246–259.
- Ellenberg U, Mattern T, Seddon PJ, Luna Jorquera G (2006) Physiological and reproductive consequences of human disturbance in Humboldt Penguins: the need for species-specific visitor management. *Biol Conserv* 133: 95–106.
- Ellenberg U, Setiawan AN, Cree A, Houston DM, Seddon PJ (2007) Elevated hormonal stress response and reduced reproductive output in Yellow-eyed Penguins exposed to unregulated tourism. *Gen Comp Endocrinol* 152: 54–63.
- Fahrig L (2003) Effects of habitat fragmentation on biodiversity. *Annu Rev Ecol Syst* 34: 487–515.
- Field A (2005) *Discovering Statistics using SPSS*. Sage Publications, London.
- Foley JA, Defries R, Asner GP, Barford C, Bonan G, Carpenter SR, Chapin FS, Coe MT, Daily GC, Gibbs HK et al. (2005) Global consequences of land use. *Science* 309: 570–574.
- Fowler GS (1999) Behavioral and hormonal responses of Magellanic Penguins (*Spheniscus magellanicus*) to tourism and nest site visitation. *Biol Conserv* 90: 143–149.
- Fox J, Weisberg S (2011) *An R Companion to Applied Regression*. Sage Publications, London.
- French SS, Denardo DF, Greives TJ, Strand CR, Demas GE (2010) Human disturbance alters endocrine and immune responses in the Galapagos marine iguana (*Amblyrhynchus cristatus*). *Horm Behav* 58: 792–799.
- Ganswindt A, Palme R, Heistermann M, Borrigan S, Hodges JK (2003) Non-invasive assessment of adrenocortical function in the male African elephant (*Loxodonta africana*) and its relation to musth. *Gen Comp Endocrinol* 134: 156–166.
- Ganzhorn JU (1995) Low-level forest disturbance effects on primary production, leaf chemistry, and lemur populations. *Ecology* 76: 2084–2096.
- Gobush KS, Mutayoba BM, Wasser SK (2008) Long-term impacts of poaching on relatedness, stress physiology, and reproductive output of adult female African elephants. *Conserv Biol* 22: 1590–1599.
- Gómez-Posada C, Roncancio N, Hincapie P, Betancourt A (2010) Densidad y composición de grupos en tres poblaciones de mono aullador rojo (*Alouatta seniculus*) en Valle y Cauca, Colombia. *Boletín Científico del Centro de Museos, Universidad de Caldas* 14: 79–91.
- González-Solis J, Guix JC, Mateos E, Llorens L (2001) Population density of primates in a large fragment of the Brazilian Atlantic rainforest. *Biodivers Conserv* 10: 1267–1282.
- Goymann W (2012) On the use of non-invasive hormone research in uncontrolled, natural environments: the problem with sex, diet, metabolic rate and the individual. *Methods Ecol Evol* 3: 757–765.
- Hannah L, Carr JL, Lankerani A (1995) Human disturbance and natural habitat: a biome level analysis of a global data set. *Biodivers Conserv* 4: 128–155.
- Harcourt AH, Coppeto SA, Parks SA (2002) Rarity, specialization and extinction in primates. *J Biogeogr* 29: 445–456.
- Heistermann M, Ademmer C, Kaumanns W (2004) Ovarian cycle and effect of social changes on adrenal and ovarian function in *Pygathrix nemaeus*. *Int J Primatol* 25: 689–708.
- Henle K, Davie KF, Kleyer M, Margules C, Settele J (2004) Predictors of species sensitivity to fragmentation. *Biodivers Conserv* 13: 207–251.
- Hinam HL, St Clair CC (2008) High levels of habitat loss and fragmentation limit reproductive success by reducing home range size and provisioning rates of Northern Saw-whet Owls. *Biol Conserv* 141: 524–535.
- Homan RN, Regosin JV, Rodrigues DM, Reed JM, Windmiller BS, Romero LM (2003) Impacts of varying habitat quality on the physiological stress of spotted salamanders (*Ambystoma maculatum*). *Anim Conserv* 6: 11–18.
- Hunt KE, Wasser SK (2003) Effect of long-term preservation methods on fecal glucocorticoid concentrations of grizzly bear and African elephant. *Physiol Biochem Zool* 76: 918–928.
- Irwin MT (2008) Diademed sifaka (*Propithecus diadema*) ranging and habitat use in continuous and fragmented forest: higher density but lower viability in fragments? *Biotropica* 40: 231–240.
- IUCN (2012) The IUCN Red List of Threatened Species. Version 2012.2. <http://www.iucnredlist.org> (last accessed 5 December 2012).
- Jaimez NA, Bribiescas RG, Aronsen GP, Anestis SA, Watts DP (2012) Urinary cortisol levels of gray-cheeked mangabeys are higher in disturbed compared to undisturbed forest areas in Kibale National Park, Uganda. *Anim Conserv* 15: 242–247.
- Janin A, Léna J-P, Joly P (2011) Beyond occurrence: body condition and stress hormone as integrative indicators of habitat availability and fragmentation in the common toad. *Biol Conserv* 144: 1008–1016.
- Johns AD (1988) Effects of “selective” timber extraction on rain forest structure and composition and some consequences for frugivores and folivores. *Biotropica* 20: 31–37.
- Johns AD, Skorupa JP (1987) Responses of rain-forest primates to habitat disturbance: a review. *Int J Primatol* 8: 157–191.
- Julliot C, Sabatier D (1993) Diet of the red howler monkey (*Alouatta seniculus*) in French Guiana. *Int J Primatol* 14: 527–550.

- Keay JM, Singh J, Gaunt MC, Kaur T (2006) Fecal glucocorticoids and their metabolites as indicators of stress in various mammalian species: a literature review. *J Zoo Wildl Med* 37: 234–244.
- Keselman HJ, Algina J, Kowalchuk RK (2001) The analysis of repeated measures designs: a review. *Br J Math Stat Psychol* 54: 1–20.
- Klein LL, Klein DJ (1977) Feeding behaviour of the Colombian spider monkey, *Ateles belzebuth*. In TH Clutton-Brock, ed., *Primate Ecology: Studies of Feeding and Ranging Behavior in Lemurs, Monkeys, and Apes*. Academic Press, London, pp 153–181.
- Landys MM, Ramenofsky M, Wingfield JC (2006) Actions of glucocorticoids at a seasonal baseline as compared to stress-related levels in the regulation of periodic life processes. *Gen Comp Endocrinol* 148: 132–149.
- Link A, de Luna AG, Alfonso F, Giraldo-Beltran P, Ramirez F (2010) Initial effects of fragmentation on the density of three neotropical primate species in two lowland forests of Colombia. *Endang Species Res* 13: 41–50.
- Lopez GO, Terborgh J, Ceballos N (2005) Food selection by a hyperdense population of red howler monkeys (*Alouatta seniculus*). *J Trop Ecol* 21: 445–450.
- Lucas JR, Freeberg TM, Egbert J, Schwabl H (2006) Fecal corticosterone, body mass, and caching rates of Carolina chickadees (*Poecile carolinensis*) from disturbed and undisturbed sites. *Horm Behav* 49: 634–643.
- Martínez-Mota R, Valdespino C, Sánchez-Ramos MA, Serio-Silva JC (2007) Effects of forest fragmentation on the physiological stress response of black howler monkeys. *Anim Conserv* 10: 374–379.
- Menon S, Poirier FE (1996) Lion-tailed macaques (*Macaca silenus*) in a disturbed forest fragment: activity patterns and time budget. *Int J Primatol* 17: 969–985.
- Michalski F, Peres CA (2005) Anthropogenic determinants of primate and carnivore local extinctions in a fragmented forest landscape of southern Amazonia. *Biol Conserv* 124: 383–396.
- Millsbaugh JJ, Washburn BE (2004) Use of fecal glucocorticoid metabolite measures in conservation biology research: considerations for application and interpretation. *Gen Comp Endocrinol* 138: 189–99.
- Milton K (1980) *The Foraging Strategy of Howler Monkeys: A Study of Primate Economics*. Columbia University Press, New York.
- Mittermeier RA, Rylands AB, Taylor LA, Chiozza F, Williamson EA, Wallis J, Schwitzer C (2012) *Primates in Peril: The World's 25 Most Endangered Primates 2012–2014*. IUCN/SSC Primate Specialist Group (PSG), International Primatological Society (IPS), Conservation International (CI), and Bristol Conservation and Science Foundation, Bristol, UK.
- Monclús R, Rödel HG, Von Holst D, De Miguel J (2005) Behavioural and physiological responses of naïve European rabbits to predator odour. *Anim Behav* 70: 753–761.
- Muehlenbein MP, Ancrenaz M, Sakong R, Ambu L, Prall S, Fuller G, Raghanti MA (2012) Ape conservation physiology: fecal glucocorticoid responses in wild *Pongo pygmaeus morio* following human visitation. *PLoS One* 7: e33357. doi:10.1371/journal.pone.0033357.
- Muller MN, Wrangham RW (2004) Dominance, cortisol and stress in wild chimpanzees (*Pan troglodytes schweinfurthii*). *Behav Ecol Sociobiol* 55: 332–340.
- Müllner A, Linsenmair EK, Wikelski M (2004) Exposure to ecotourism reduces survival and affects stress response in hoatzin chicks (*Opisthocomus hoazin*). *Biol Conserv* 118: 549–558.
- Munshi-South J, Tchignoumba L, Brown J, Abbondanza N, Maldonado JE, Henderson A, Alonso A (2008) Physiological indicators of stress in African forest elephants (*Loxodonta africana cyclotis*) in relation to petroleum operations in Gabon, Central Africa. *Divers Distrib* 14: 995–1003.
- Neville MK (1972) The population structure of red howler monkeys (*Alouatta seniculus*) in Trinidad and Venezuela. *Folia Primatol* 17: 56–86.
- Ostro LET, Silver SC, Koontz FW, Horwich RH, Brockett R (2001) Shifts in social structure of black howler (*Alouatta pigra*) groups associated with natural and experimental variation in population density. *Int J Primatol* 22: 733–748.
- Peres CA (2001) Synergistic effects of subsistence hunting and habitat fragmentation on Amazonian forest vertebrates. *Conserv Biol* 15: 1490–1505.
- Pickering AD, Pottinger TG, Sumpter JP, Carragher JF, Le Bail PY (1991) Effects of acute and chronic stress on the levels of circulating growth hormone in the rainbow trout, *Oncorhynchus mykiss*. *Gen Comp Endocrinol* 83: 86–93.
- Piñeiro A, Barja I, Silván G, Illera JC (2012) Effects of tourist pressure and reproduction on physiological stress response in wildcats: management implications for species conservation. *Wildl Res* 39: 532–539.
- Purvis A, Gittleman JL, Cowlishaw G, Mace GM (2000) Predicting extinction risk in declining species. *Proc Biol Sci* 267: 1947–1952.
- Putz FE, Leigh EGJ, Wright SJ (1990) Solitary confinement in Panama. *Garden* 2: 18–23.
- R Development Core Team (2012) *A language and environment for statistical computing*. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org>
- Rangel-Negrín A, Alfaro JL, Valdez RA, Romano MC, Serio-Silva JC (2009) Stress in Yucatan spider monkeys: effects of environmental conditions on fecal cortisol levels in wild and captive populations. *Anim Conserv* 12: 496–502.
- Rimbach R, Heymann EW, Link A, Heistermann M (2013) Validation of an enzyme immunoassay for assessing adrenocortical activity and characterization of factors that affect fecal glucocorticoid metabolite levels in two New World primates. *Gen Comp Endocrinol* 191: 13–23.
- Rumiz DI (1990) *Alouatta caraya*: population density and demography. *Am J Primatol* 21: 279–294.
- Sapolsky RM, Romero ML, Munck AU (2000) How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocr Rev* 21: 55–89.
- Selye H (1956) *The Stress of Life*. McGraw-Hill, New York.

- Setchell JM, Smith T, Wickings EJ, Knapp LA (2010) Stress, social behaviour, and secondary sexual traits in a male primate. *Horm Behav* 58: 720–728.
- Shutt K, Setchell JM, Heistermann M (2012) Non-invasive monitoring of physiological stress in the Western lowland gorilla (*Gorilla gorilla gorilla*): validation of a fecal glucocorticoid assay and methods for practical application in the field. *Gen Comp Endocrinol* 179: 167–177.
- Strasser EH, Heath JA (2013) Reproductive failure of a human-tolerant species, the American kestrel, is associated with stress and human disturbance. *J Appl Ecol* 50: 912–919.
- Stratakis CA, Chrousos GP (1995) Neuroendocrinology and pathophysiology of the stress system. *Ann NY Acad Sci* 77: 1–18.
- Sumner J, Moritz C, Shine R (1999) Shrinking forest shrinks skink: morphological change in response to rainforest fragmentation in the prickly forest skink (*Gnypetoscincus queenslandiae*). *Biol Conserv* 91: 159–167.
- Symington MM (1988) Food competition and foraging party size in the black spider monkey (*Ateles paniscus chamek*). *Behaviour* 105: 117–134.
- Tecot SR (2008) Seasonality and predictability: the hormonal and behavioral responses of the red-bellied lemur, *Eulemur rubriventer*, in southeastern Madagascar. PhD thesis. University of Texas at Austin.
- Thiel D, Palme R, Jenni L, Jenni-Eiermann S (2011) Winter tourism increases stress hormone levels in the Capercaillie *Tetrao urogallus*. *Ibis* 153: 122–133.
- Umapathy G, Hussain S, Shivaji S (2011) Impact of habitat fragmentation on the demography of lion-tailed macaque (*Macaca silenus*) populations in the rainforest of Anamalai Hills, Western Ghats, India. *Int J Primatol* 32: 889–900.
- Urbani B, Morales AL, Link A, Stevenson PR (2008) *Ateles hybridus*. In: IUCN 2013. IUCN Red List of Threatened Species. Version 2013.1. <http://www.iucnredlist.org>. (last accessed 25 July 2013).
- von der Ohe CG, Wasser SK, Hunt KE, Servheen C (2004) Factors associated with fecal glucocorticoids in Alaskan brown bears (*Ursus arctos horribilis*). *Physiol Biochem Zool* 77: 313–320.
- Wasser SK, Bevis K, King G, Hanson E (1997) Noninvasive physiological measures of disturbance in the Northern Spotted Owl. *Conserv Biol* 11: 1019–1022.
- Wasser SK, Davenport B, Ramage ER, Hunt KE, Parker M, Clarke C, Stenhouse G (2004) Scat detection dogs in wildlife research and management: application to grizzly and black bears in the Yellowhead Ecosystem, Alberta, Canada. *Can J Zool* 82: 475–492.
- Wingfield JC, Maney DL, Breuner CW, Jacobs JD, Lynn S, Ramenofsky M, Richardson RD (1998) Ecological bases of hormone-behavior interactions: the “emergency life history stage”. *Integr Comp Biol* 38: 191–206.
- Zwijacz-Kozica T, Selva N, Barja I, Silván G, Martínez-Fernández L, Illera JC, Jodłowski M (2012) Concentration of fecal cortisol metabolites in chamois in relation to tourist pressure in Tatra National Park (South Poland). *Acta Theriol* 58: 215–222.